General information for administering compresses

Compress materials

Cloth sizes that are used for all large body parts that have proven themselves:

- Inner cloth = substance cloth (smooth cotton, double layered) width approx. 28 cm
- Middle cloth (rough cotton, double layered, or terry cloth) width approx. 32 cm
- Outer cloth (wool or molleton fabric, or terry cloth towel) width approx. 36 cm
- A length of 140 cm is the same for all cloths.

Children's compresses have a length of one and a half times their upper body circumference.

The width is measured from the waist to the armpit.

For **oil compresses** and **poultices** you can use old cotton cloths (old handkerchiefs, tea towels, sheets torn to size).

Preparing the patient, room and materials Patient

- At least half an hour after the last meal
- Empty the bladder before the treatment
- The feet must be warm! If necessary, warm them up with a hot water bottle or foot bath (otherwise there is a danger that the warmth coming from the compress will rise towards the head and the feet will become even colder).
- Remove all confining clothing parts, including watches, jewelry, etc.

Room Good ventilation, pleasantly warm, no draughts; no background noise; no direct incidence of light on the patient's eyes, this also applies to the post-treatment rest (see below).

Materials Everything that is needed for the application must be prepared beforehand and placed on the bed ready to use: knee roll, small pillow for the neck, compress utensils.

Hot-water bottle: filled (not bulging) with hot water, vented (= place the hot-water bottle flat on the table, hold the filler neck upright so that all the air can escape).

Positioning the patient and applying the compress

If possible, have the patient lie on their back. If necessary, elevate their upper body and support their arms. Lay out the outer and middle cloths on the bed at the height of the corresponding part of the body, have the patient lie on them, mold the cloths one after the other over the substance cloth.

Place the knee roll or similar, cover the patient, including their shoulders. Avoid constriction in the neck area. If the person feels constricted in the heart area and/or breathless, leave their hands uncovered.

Place a small pillow under their neck so that they do not have to hold their head themselves. Remove the hot-water bottle from their feet! Cover their feet, even from the sides, without putting pressure on their toes. Do not pull the blanket under their heels.

Temperature Hot moist compresses: fan the body part with the hot cloth, then apply as hot as possible. Fevers from approx. 39°C: Choose a compress solution 2°C cooler than the body temperature; this has a cooling effect.

Preparing tea Pour 1 liter boiling water over 1 tablespoon of dried leaves,

let steep for 3–5 minutes and strain into a thermos flask.

Essential oils 1 teaspoon in a cup of water or **1** tablespoon in approx. ½ liter of water

Post-treatment rest Every compress needs a rest afterwards. This is an essential part of the application and starts with the removal of the compress materials. As a rule, it is 30 minutes.

Follow-up After removing the cloths, rinse the inner cloth in clear water and hang them all up to dry. They can be reused several times. Poultice packs are only used once.

Instructions: Onion Ear Bags

Materials

- 1 onion
- Cloth handkerchief as a substance cloth
- Cap to fix the compress in place
- Knife and cutting board
- Freezer bag
- Hot-water bottle, filled hot (70°C or 158°F), de-aired

Instructions

- Peel the onion and chop into small pieces
- Spread the onion pieces over an area of approx. 7 x 7 cm on the middle of the unfolded handkerchief
- Fold the handkerchief into a small pack
- Place the pack in the freezer bag
- Place the single-layer fabric side of the pack on the hot-water bottle for a few minutes.
- Remove the heated onion pack from the freezer bag
- Place the pack on the affected ear and place it around the auricle so that the area behind the ear is also covered
- Fix the bag in place with the cap (alternatively you can use a scarf, but a cap holds it better in place)
- If possible, leave the compress on for 2 hours

Follow-up

- Discard the onion
- Wash out the handkerchief
- If necessary, wash off the neck with warm water to get rid of the onion smell

Evidence

Well-proven in many patients

Dosage

1 x daily

Onset of effect

Rapid pain alleviation

Length of therapy

As required, often a single application is all that is needed

[©] Vademecum of External Applications 12/2019; www.vademecum.org; may only be duplicated and passed on to others if the source is named.